This report contains data through the week ending 1/11/2014 (MMWR week 2).



Overview of Influenza Surveillance: Surveillance for the 2013-2014 influenza season officially began on September 29, 2013. The Utah Department of Health publishes a weekly report throughout the active influenza season that synthesizes data from a variety of sources to give the most complete and up-to-date picture of influenza activity in the state of Utah. Data in this report should be considered provisional, and may change as more complete reports are received.

Influenza-like Illness (ILI): The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet) is a national system that conducts surveillance for influenza-like illness (ILI) in outpatient healthcare facilities. ILINet providers report weekly the total number of patients seen for any reason and the number of patients seen with ILI (defined as a fever ≥ 100° F and a cough or sore throat). These data are used to determine the amount of ILI circulating in the community, as well as provide insight into regional differences in ILI activity. Currently, more than 50 facilities throughout Utah participate in ILINet.




Table 1. Influenza-like Illness (ILI) Activity Levels by Health District - Utah, Current Week

| Ticaltii District | Otall, Callelle Week |
|------------------------|----------------------|
| Health District | ILI Activity |
| Bear River | Low/Moderate |
| Central | Low/Moderate |
| Davis | Low/Moderate |
| Salt Lake | Low/Moderate |
| Southeastern | No Data |
| Southwest | Low/Moderate |
| Summit | Minimal |
| Tooele | Minimal |
| TriCounty | No Data |
| Utah | Low/Moderate |
| Wasatch | Minimal |
| Weber-Morgan | Low/Moderate |
| State | Low/Moderate |

This report contains data through the week ending 1/11/2014 (MMWR week 2).



Influenza Hospitalizations: Influenza hospitalizations are a reportable condition in Utah. A person meets the case definition for an influenza hospitalization if they are hospitalized for any length of time and have an influenza positive serology, DFA, PCR, culture or rapid influenza diagnostic test. Public health in Utah gathers a variety of data on influenza hospitalizations including clinical features, course of illness, risk and protective factors, and influenza type and subtype. Data from influenza hospitalizations allows public health in Utah to better understand subgroups of the Utah population that are most severely affected by influenza and help to guide prevention messages and interventions.




Table 2. Influenza Hospitalizations by Health District - Utah

| Health District | Current Week | Season To Date |
|------------------------|--------------|----------------|
| Bear River | 1 | 13 |
| Central | 2 | 7 |
| Davis | 4 | 31 |
| Salt Lake | 22 | 264 |
| Southeastern | 0 | 1 |
| Southwest | 7 | 32 |
| Summit | 1 | 5 |
| Tooele | 0 | 5 |
| TriCounty | 1 | 5 |
| Utah | 3 | 55 |
| Wasatch | 2 | 5 |
| Weber-Morgan | 6 | 55 |
| State | 49 | 478 |

This report contains data through the week ending 1/11/2014 (MMWR week 2).



Table 3. Influenza Hospitalizations by Age Group - Utah, Season To Date


| Age Group | Total Cases | % of Cases | Rate* |
|-----------|-------------|------------|-------|
| 0-4 | 60 | 12.6 | 22.58 |
| 5-24 | 43 | 9.0 | 4.49 |
| 25-49 | 155 | 32.4 | 16.24 |
| 50-64 | 127 | 26.6 | 30.67 |
| 65+ | 93 | 19.5 | 35.41 |
| Total | 478 | 100.0 | 16.74 |

^{*}Rate is calculated as the number of cases per 100,000 population

Table 4. Influenza Hospitalizations by Sex and Race - Utah, Season To Date

| Variab | le | Num. of Cases | % of Cases | % in Utah Pop p value* |
|--------|----------------------------------|---------------|------------|------------------------|
| Sex | Male | 259 | 54.2 | 50.2 0.0851 |
| | Female | 219 | 45.8 | 49.8 0.0847 |
| | Unknown | 0 | 0.0 | NA |
| Race | White, Not Hispanic | 315 | 65.9 | 79.9 < 0.0001 |
| | Hispanic | 77 | 16.1 | 11.9 0.0004 |
| | Native Hawaiian/Pacific Islander | 27 | 5.6 | 1.0 < 0.0001 |
| | Black/African American | 8 | 1.7 | 1.3 0.3498 |
| | American Indian | 5 | 1.0 | 1.5 0.5177 |
| | Asian | 2 | 0.4 | 2.2 0.0120 |
| | Unknown | 44 | 9.2 | NA |

^{*}If a p value is \leq 0.05, there is a significant difference between the percentage seen in influenza hospitalizations and the general Utah population.



This report contains data through the week ending 1/11/2014 (MMWR week 2).



Student Absenteeism: School-age children are at high risk for respiratory virus infections, including influenza. Aggregate, all-cause absenteeism data is collected weekly from over 350 schools throughout Utah. These data are analyzed to identify elevated absenteeism rates that could indicate the circulation of influenza in school-age children.





Table 6. Weekly Student Absenteeism - Utah, Current Week

| Health District | Absences per 100 students/day |
|-----------------|-------------------------------|
| Bear River | 3.6 |
| Central | 4.6 |
| Davis | 4.0 |
| Salt Lake | 3.8 |
| Southeast | 5.2 |
| Southwest | 5.2 |
| Summit | 5.7 |
| Tooele | 5.2 |
| TriCounty | 6.5 |
| Utah | 1.9 |
| Wasatch | 4.2 |
| Weber-Morgan | 4.8 |
| State | 4.3 |


Pneumonia and Influenza Deaths: Each week the total number of death certificates received and the number of those for which pneumonia or influenza was listed as an underlying or contributing cause of death is collected. The percentage of deaths due to pneumonia and influenza are compared with a seasonal baseline and epidemic threshold value calculated for each week. These data are used to monitor the severity of influenza illness in the community.



This report contains data through the week ending 1/11/2014 (MMWR week 2).



Laboratory Surveillance: The Utah Public Health Laboratory recieves specimens from all over the state for comprehensive influenza testing. All specimens are tested to determine influenza type and subtype. A portion of specimens are also sent to the Centers for Disease Control and Prevention for additional testing, including gene sequencing, antiviral resistance testing and antigenic characterization.



Note: Due to limited resources, testing at UPHL has been restricted to a subset of the submitted specimens since the week of 12/7/2013. The decrease in positive specimens in the graph above represents the change in testing practices, and not a decrease in circulating virus.

Table 8. Utah Public Health Laboratory Influenza Testing Data

| | Current Week | | Season ⁻ | Γο Date | |
|------------------------------------|--------------|---------|---------------------|---------|--|
| | Total | Percent | Total | Percent | |
| Specimens tested | 59 | | 552 | | |
| Positive specimens | 20 | 33.9 | 256 | 46.4 | |
| Positive Specimens by Type/Subtype | | | | | |
| Influenza A | 20 | 100.0 | 252 | 98.4 | |
| A (2009 H1N1) | 20 | 100.0 | 248 | 98.4 | |
| A (H1) | 0 | 0.0 | 0 | 0.0 | |
| A (H3) | 0 | 0.0 | 4 | 1.6 | |
| A (unable to subtype) | 0 | 0.0 | 0 | 0.0 | |
| Influenza B | 0 | 0.0 | 4 | 1.6 | |